



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).



$$\therefore a \cos \frac{ns}{a} = x + \frac{\sqrt{a^2 \left[ 1 + \left( \frac{dy}{dx} \right)^2 \right] - \left( y - x \frac{dy}{dx} \right)^2 - \left( x + y \frac{dy}{dx} \right)^2}}{1 + \left( \frac{dy}{dx} \right)^2}.$$

$$\text{If } CR = y_1, ns = a \sin^{-1} \frac{y_1}{a} = a \sin^{-1} \left( \frac{y + CR'}{a} \right),$$

$$\text{but } CR' = m \sin \phi = m \frac{dy}{ds}. \quad \therefore a \sin \frac{ns}{a} = y + m \frac{dy}{ds} \dots (2).$$

$$\therefore a \sin \frac{ns}{a} = y + \frac{\sqrt{a^2 \left\{ 1 + \left( \frac{dy}{dx} \right)^2 \right\} - \left( y + x \frac{dy}{dx} \right)^2 - \left( x + y \frac{dy}{dx} \right)^2}}{\sqrt{\left\{ 1 + \left( \frac{dy}{dx} \right)^2 \right\} \left\{ 1 + \left( \frac{dx}{dy} \right)^2 \right\}}}.$$

The solution of either of these differential equations (if possible) give the rectangular equation to the path of the pursuer. After we know the equation to the curve we can find its length, from which we know the time and distance. If the pursuer's velocity is less than that of the pursued the race will last an infinitely long time, or the *pursued* will catch the *pursuer* and thus end the race.

II. Solution by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

Let  $O$  and  $B$  represent the starting-points of  $A$  and  $B$ , then will  $P$  and  $C$  represent their positions at any time after starting. Make  $OB = R = 15$  feet,  $n = v \div u = 6$ ,  $\angle BOC = \psi$ ,  $\angle BOG = \theta$ , and  $BQC = \phi$ . Obviously the polar and the Cartesian co-ordinates of the point  $P$  located on the required curve, are respectively  $(r, \theta)$  and  $(x, y)$ . Hence  $dy \div dx = \tan \phi$ ,

$$OQ = y(dx \div dy) = y \cot \phi, \angle OPQ = (\phi - \theta), \text{ and } POC = (\psi - \theta).$$

$$\text{Now } PQ = \left( \frac{\sin(\psi - \theta)}{\sin(\phi - \theta)} \right) OQ = y \operatorname{cosec} \phi \dots (1). \text{ That is, } \frac{\sin(\psi - \theta)}{\sin(\phi - \theta)}$$

$$= \tan \phi \operatorname{cosec} \phi \dots (2).$$

$$\therefore \sin \psi$$

$$= \frac{-(\tan \phi - \tan \theta) \pm \sqrt{(\tan \phi - \tan \theta)^2 + (1 + \tan^2 \theta)[2 \tan \phi \tan \theta - \tan^2 \phi]}}{1 + \tan^2 \theta};$$

$$\text{and } \psi = \sin^{-1} \left( \frac{(\tan \phi - \tan \theta) \pm \tan \theta \sqrt{(1 + 2 \tan \phi \tan \theta - \tan^2 \phi)}}{1 + \tan^2 \theta} \right),$$

$$= \sin^{-1} \left( \frac{(\tan \phi - \tan \theta) \pm \tan \theta \sqrt{[(1 + \tan^2 \theta) - (\tan \phi - \tan \theta)^2]}}{1 + \tan^2 \theta} \right).$$

$$\text{But } s = \int \sqrt{1 + \left( \frac{dy}{dx} \right)^2} dx; \tan \theta = y \div x; \text{ and from the problem, } ns = R \dots (3).$$

$$\text{By obvious transformations, we have from (3) } \left( 1 + \frac{y^2}{x^2} \right) \sin \left( \frac{ns}{R} \right) =$$

$$\left\{ \left( \frac{dy}{dx} - \frac{y}{x} \right) \pm \frac{y}{x} \sqrt{\left[ \left( 1 + \frac{y^2}{x^2} \right) - \left( \frac{dy}{dx} - \frac{y}{x} \right)^2 \right]} \right\}, \text{ which is the Cartesian differen-}$$

tial equation of the required curve; and this equation does not appear to be integrable. Many other differential equations of the required curve can be deduced; but all of these equations, as to their integrability, transcend the present limits of mathematical genius.

III. Solution By JAMES McHAHON, M. A., Associate Editor of the "Annals of Mathematics", Department of Mathematics, Cornell University, Ithaca, New York.

Let arc  $BC = \nu$ ,  $\angle DCQ = \mu$ ,  $P, P'$  two consecutive points on the curve, then  $dm = C'P' - CP = C'A - CA - (AP + AP') = CC' \cos \mu - \frac{u}{v} CC'$ .

Let  $\frac{u}{v} = n'$ .  $\therefore dm = d\nu(\cos \mu - n') \dots (3)$ .  $d\mu = C'OC - CAC = \frac{d\nu}{a} - \frac{d\nu \sin \mu}{m}$ .

$\therefore am d\mu = (m - a \sin \mu) d\nu \dots (4)$ . From (3) and (4) by elimination and solution (if possible) the equation to the curve of pursuit is found.

IV. Comment by M. C. STEVENS, M. A., Professor of Mathematics, Purdue University, Lafayette, Indiana.

To solve this problem, if we let  $(x', y')$  be the co-ordinates of a point of the pursued at any moment, and  $(x, y)$  the coordinates of the pursuer at the corresponding moment, then we have,  $x'^2 + y'^2 = 225 \dots (1)$ .  $y' - y = \frac{dy}{dx}(x' - x) \dots (2)$

$$\sqrt{\left(\frac{dx'}{dx}\right)^2 + \left(\frac{dy'}{dx}\right)^2} = 6\sqrt{1 + \left(\frac{dy}{dx}\right)^2} \dots (3)$$

By elimination we can find a differential equation involving,  $x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}$ . (See Boole's Diff. Equations, page 251).

But this results in a very complicated equation, which has never, so far as I know, been solved. Now since the velocity of  $B$  is greater than that of  $A$ ,  $A$  will never overtake  $B$ ; hence the answer to the question is, the *time* is *infinite* and the *distance* is *infinite*. (See remarks on Curve of Pursuit in Runkle's Mathematical Monthly, Vol. I, p. 248).

## PROBLEMS.

25. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy, New Windsor College, New Windsor, Maryland.

The leaf of the curve: "The Devil on Two Sticks", equation  $y^4 - x^4 + 100a^2x^2 - 96a^2y^2 = 0$ , revolves around the axis of  $x$ . Deduce the expression for the volume generated.

26. Proposed by Professor J. F. W. SCHEFFER, M. A., Hagerstown, Maryland.

According to Bessel the ratio of the squares of the polar diameter of the earth to that of the equatorial diameter is .9933254. Find at what altitude the angle made by a body falling to the earth with a perpendicular to the surface is greatest. Find also this maximum angle.

Solutions to these problems should be received on or before November 1st.